

Table 27: Percentage of caretakers of children 0-59 months who know at least 2 signs for seeking care immediately, Country, Year

	Knows child should be taken to health facility if child:							Knows at least two signs	Number of caretakers
	Not able to drink /breastfeed	Becomes sicker	Develops a fever	Has fast breathing	Has difficult breathing	Has blood in stool	Is drinking poorly		
Central	.0	8.7	68.0	.9	6.9	1.1	.6	10.3	663
South Central	.3	15.2	68.7	2.6	8.1	4.5	3.7	17.6	370
West	10.4	16.1	74.0	13.3	17.2	11.4	11.4	21.9	270
East	.9	10.1	64.5	2.6	5.3	.9	.4	9.6	219
South	14.0	22.4	91.6	21.1	24.2	14.6	14.0	34.2	309
Urban	.4	10.2	69.0	1.4	6.3	1.5	1.2	11.2	896
Rural	7.6	16.8	76.1	11.8	16.3	9.4	8.8	23.4	934
Mother's education									
None	6.9	19.1	83.1	11.3	17.4	7.2	7.7	26.8	238
Primary	7.7	17.1	74.0	11.4	16.0	9.7	8.3	20.1	496
Secondary +	1.8	10.8	69.7	3.6	8.0	3.3	3.1	14.2	1096
Total	4.0	13.6	72.6	6.7	11.4	5.6	5.1	17.4	1830